

# The Translator.



## Computes all.

All there is to compute. Directly onto tape. Blue Shield enrollment information. Medicare information. Information information. The Translator means time saved in transcription. Time saved in key punch. Time saved for you. So in 1970, you'll have more time to be a doctor.



GREATER NEW YORK'S

**BLUE SHIELD**

UNITED MEDICAL SERVICE, INC.

## Service is what we're all about.

The Bulletin of the New York Academy of Medicine, Vol. 46, No. 5, May 1970. Published monthly by the New York Academy of Medicine, 2 East 103 Street, New York, N. Y. 10029. Entered as second-class matter February 3, 1928, at the Post Office at New York, N. Y., under the act of August 24, 1912. Postage paid at New York, N. Y. Annual subscription United States \$15.00. Canada \$16.00. All other countries \$17.00. Single copies \$3.00.

# 3

Polymyxin B—Bacitracin—Neomycin

# against 10




Pseudomonas




Hemophilus




Klebsiella



Aerobacter




Escherichia



Proteus




Corynebacterium



Staphylococcus



Streptococcus



Pneumococcus

## ‘Neosporin’<sup>®</sup> Ointment Polymyxin B—Bacitracin—Neomycin

Overlapping, broad bactericidal coverage.

Nonirritant ointment base; also enhances spreading and penetration.

Each gram contains:

‘Aerosporin’<sup>®</sup>

brand Polymyxin B Sulfate . . . . . 5,000 Units

Zinc Bacitracin . . . . . 400 Units

Neomycin Sulfate . . . . . 5 mg.

(equivalent to 3.5 mg. Neomycin Base)

Special White Petrolatum . . . . . q.s.

**Contraindications:** This product is contraindicated in those individuals who have shown hypersensitivity to any of its components. Do not use in the external ear canal if the eardrum is perforated.

**Precautions:** As with other antibiotic products,

prolonged use may result in overgrowth of non-susceptible organisms, including fungi. Appropriate measures should be taken if this occurs. Articles in the current medical literature indicate an increase in the prevalence of persons allergic to neomycin. The possibility of such a reaction should be borne in mind.

**Available:** Tubes of 1 oz., 1/2 oz. with applicator tip, 1/8 oz. with ophthalmic tip. The ointment base and the formula of the various sizes are identical, but only the 1/8 oz. tube should be used for ophthalmic purposes.



**BURROUGHS WELLCOME & CO. (U.S.A.) INC., Tuckahoe, New York**

"My heart beats louder at night..."



The chronically anxious cardiac patient who becomes overly preoccupied with his condition may tend to become even more apprehensive during the nighttime hours. With senses sharpened by acute anxiety, he may interpret as "ominous" changes in his heart's action. And with intensified anxiety, sleep may often be disturbingly elusive.

To help alleviate anxiety-induced insomnia, an added h.s. dose of Librium® (chlordiazepoxide HCl) can be particularly effective. Through its prompt, dependable anxiolytic action, Librium frequently calms the patient and encourages relaxation and sleep. A t.i.d. daytime regimen often helps to decrease excessive "cardiac anxiety" in conjunction with the physician's reassurance and counsel, enabling the patient to adopt a more realistic attitude toward his disorder and toward therapy. Also of therapeutic importance: on proper maintenance dosage, Librium seldom impairs ability to function. In general use, the most common side effects reported have been drowsiness, ataxia and confusion, particularly in the elderly and debilitated. (See full prescribing information.)

**For anxiety-induced insomnia in the cardiac patient adjunctive**

**Librium®**  
**chlordiazepoxide HCl)**  
**t.i.d. plus h.s.**  
**5-mg, 10-mg and 25-mg capsules**



**Roche**  
**LABORATORIES**

Division of Hoffmann-La Roche Inc.  
Nutley, New Jersey 07110

Before prescribing, please consult complete product information, a summary of which follows:

**Indications:** Indicated when anxiety, tension and apprehension are significant components of the clinical profile.


**Contraindications:** Patients with known hypersensitivity to the drug.

**Warnings:** Caution patients about possible combined effects with alcohol and other CNS depressants. As with all CNS-acting drugs, caution patients against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving). Though physical and psychological dependence have rarely been reported on recommended doses, use caution in administering to addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions), following discontinuation of the drug and similar to those seen with barbiturates, have been reported. Use of any drug in pregnancy, lactation, or in women of childbearing age requires that its potential benefits be weighed against its possible hazards.

**Precautions:** In the elderly and debilitated, and in children over six, limit to smallest effective dosage (initially 10 mg or less per day) to preclude ataxia or oversedation, increasing gradually as needed and tolerated. Not recommended in children under six. Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider individual pharmacologic effects, particularly in use of potentiating drugs such as MAO inhibitors and phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function. Paradoxical reactions (e.g., excitement, stimulation and acute rage) have been reported in psychiatric patients and hyperactive aggressive children. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation have been reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship has not been established clinically.

**Adverse Reactions:** Drowsiness, ataxia and confusion may occur, especially in the elderly and debilitated. These are reversible in most instances by proper dosage adjustment, but are also occasionally observed at the lower dosage ranges. In a few instances syncope has been reported. Also encountered are isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent and generally controlled with dosage reduction; changes in EEG patterns (low-voltage fast activity) may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice and hepatic dysfunction have been reported occasionally, making periodic blood counts and liver function tests advisable during protracted therapy.

**Now  
available for your  
prescribing  
needs**




**Cordran<sup>®</sup> Tape**  
**Flurandrenolide Tape** (4 mcg. per sq. cm.)

Additional information available upon request • Eli Lilly and Company, Indianapolis, Indiana 46206

000108

*Lilly*

**now for your non-hospitalized patients**



Serum Chemistry Graph provided is 8" x 11"

# blood chemistry profiles

run on

**TECHNICON®**

Sequential Multiple Analyzer

**only \$15<sup>00</sup>**

FOR A ~~12~~-TEST BLOOD CHEMISTRY PROFILE

Heretofore available only to patients in a few large hospitals, PATHOLOGICAL LABORATORY now provides this rapid automated analytical service for the non-hospitalized patient.

A Serum Chemistry Graph of each analysis summarizes in graphic form the results of all 12 tests together, superimposed over a background of normal values.

Patients may be directed to the laboratory; or blood samples can be mailed in post-paid containers provided by our laboratory. The Serum Chemistry Graph and a report in the usual numerical format will be mailed the same day samples are received. Free messenger pick-up is available in select areas.

## PATHOLOGICAL LABORATORY

40-03 NATIONAL ST., CORONA, N.Y. 11368

TELEPHONE 212-639-5403

**send for  
post-paid  
blood sample  
mailing  
containers**

PATHOLOGICAL LABORATORY • 40-03 National St., Corona, N.Y. 11368

Gentlemen:

- ☐ Please send additional literature
- ☐ Please send post-paid containers for blood samples

Dr. ....

Street .....

City ..... State ..... Zip .....

---

---

## STUDIES IN MEDICAL GEOGRAPHY

*Edited by Jacques M. May, M.D.*

**Vol. 2. STUDIES IN DISEASE ECOLOGY**  
xx, 613 pages, tables, 14 illus. 1961 \$18.50

**Vol. 3. THE ECOLOGY OF MALNUTRITION IN THE FAR AND NEAR EAST**  
704 pages, 220 tables, 30 illus. 1961 \$13.50

**Vol. 4. THE ECOLOGY OF MALNUTRITION IN FIVE COUNTRIES OF EASTERN AND CENTRAL EUROPE: East Germany, Poland, Yugoslavia, Albania, Greece**  
xviii, 292 pages, 103 plates, 9 maps. 1964 \$10.50

**Vol. 5. THE ECOLOGY OF MALNUTRITION IN MIDDLE AFRICA: Ghana, Nigeria, Republic of the Congo, Rwanda and Burundi and the former French Equatorial Africa**  
255 pages, maps. 1965 \$9.50

**Vol. 6. THE ECOLOGY OF MALNUTRITION IN CENTRAL AND SOUTHEASTERN EUROPE: Austria, Hungary, Rumania, Bulgaria, and Czechoslovakia**  
xv, 271 pages and index, 19 maps, 91 tables. 1966 \$10.50

**Vol. 7. THE ECOLOGY OF MALNUTRITION IN NORTH-EAST AFRICA: Lybia, Tunisia, Algeria, Morocco, Spanish Sahara and Ifni, Mauretania**  
xv, 275 pages, maps, tables. 1967 \$11.50

**Vol. 8. THE ECOLOGY OF MALNUTRITION IN THE FRENCH-SPEAKING COUNTRIES OF WEST AFRICA AND MADAGASCAR: Senegal, Guinea, Ivory Coast, Togo, Dahomey, Cameroon, Niger, Mali, Upper Volta, and Madagascar**  
xiii, 433 pages, index, maps, tables. 1968. \$14.50

**Vol. 9. THE ECOLOGY OF MALNUTRITION IN FOURTEEN COUNTRIES OF TROPICAL AFRICA**  
In preparation

**HAFNER PUBLISHING COMPANY, INC.**

**P. O. Box 2000, 260 Heights Road, Darien, Conn. 06820**

---

---


**Over 5 million prescriptions already filled at Macy's**

*All 13 Macy Prescription Centres  
satisfy your patients with  
efficient professional service*



- You may phone in your patients' prescriptions (except for narcotics, amphetamines, and barbiturates).
- Your patients may charge their prescriptions on their Macy Shopping Accounts or C-T Plans.
- Your patients may have their prescriptions delivered anywhere free of extra charge.
- We serve your patient with a full line of convalescent and diabetic needs.

**Macy's Herald Square, OX 5-4400**  
**Macy's Parkchester, TA 8-7000**  
**Macy's Jamaica, OL 7-9000**  
**Macy's Flatbush, UL 6-5000**  
**Macy's Roosevelt Field, PI 6-8200**  
**Macy's Huntington, AR 1-3000**  
**Macy's White Plains, WH 6-5015**  
**Macy's New Haven, Conn. 624-9271**  
**Macy's Bay Shore, MO 5-8400**  
**Macy's Queens, AR 1-9100**  
**Macy's Colonie, 459-1950**  
**Macy's New Rochelle, 633-7700**  
**Macy's Smith Haven, 724-4900**



# When disease is ruled out and psychic tension is implicated

## Valium® (diazepam)

### helps relax the patient and relieve his somatic symptoms

Before prescribing, please consult complete product information, a summary of which follows:

**Indications:** Tension and anxiety states; somatic complaints which are concomitants of emotional factors; psychoneurotic states manifested by tension, anxiety, apprehension, fatigue, depressive symptoms or agitation; acute agitation, tremor, delirium tremens and hallucinosis due to acute alcohol withdrawal; adjunctively in skeletal muscle spasm due to reflex spasm to local pathology, spasticity caused by upper motor neuron disorders, athetosis, stiff-man syndrome, convulsive disorders (not for sole therapy).

**Contraindicated:** Known hypersensitivity to the drug. Children under 6 months of age. Acute narrow angle glaucoma.

**Warnings:** Not of value in psychotic patients. Caution against hazardous occupations requiring complete mental alertness. When used adjunctively in convulsive disorders, possibility of increase in frequency and/or severity of grand mal seizures may require increased dosage of standard anticonvulsant medication; abrupt withdrawal may be associated with temporary increase in frequency and/or severity of seizures. Advise against simultaneous ingestion of alcohol and other CNS depressants. Withdrawal symptoms have occurred following abrupt discontinuance. Keep addiction-prone individuals under careful surveillance because of their predisposition to habituation and dependence. In pregnancy, lactation

or women of childbearing age, weigh potential benefit against possible hazard.

**Precautions:** If combined with other psychotropics or anticonvulsants, consider carefully pharmacology of agents employed. Usual precautions indicated in patients severely depressed, or with latent depression, or with suicidal tendencies. Observe usual precautions in impaired renal or hepatic function. Limit dosage to smallest effective amount in elderly and debilitated to preclude ataxia or oversedation.

**Side Effects:** Drowsiness, confusion, diplopia, hypotension, changes in libido, nausea, fatigue, depression, dysarthria, jaundice, skin rash, ataxia, constipation, headache, incontinence, changes in salivation, slurred speech, tremor, vertigo, urinary retention, blurred vision. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances, stimulation, have been reported; should these occur, discontinue drug. Isolated reports of neutropenia, jaundice; periodic blood counts and liver function tests advisable during long-term therapy.



**Roche**  
LABORATORIES  
Division of Hoffmann-La Roche Inc.  
Nutley, New Jersey 07110